

Are General Quantum Correlations Monogamous?

Alexander Streltsov,¹ Gerardo Adesso,² Marco Piani,³ and Dagmar Bruß¹

¹*Heinrich-Heine-Universität Düsseldorf, Institut für Theoretische Physik III, D-40225 Düsseldorf, Germany*

²*School of Mathematical Sciences, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom*

³*Institute for Quantum Computing and Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada*

(Received 16 December 2011; published 1 August 2012)

Quantum entanglement and quantum nonlocality are known to exhibit monogamy; that is, they obey strong constraints on how they can be distributed among multipartite systems. Quantum correlations that comprise and go beyond entanglement are quantified by, e.g., quantum discord. It was observed recently that for some states quantum discord is not monogamous. We prove, in general, that any measure of correlations that is monogamous for all states and satisfies reasonable basic properties must vanish for all separable states: only entanglement measures can be strictly monogamous. Monogamy of other than entanglement measures can still be satisfied for special, restricted cases: we prove that the geometric measure of discord satisfies the monogamy inequality on all pure states of three qubits.

DOI: 10.1103/PhysRevLett.109.050503

PACS numbers: 03.67.-a

Entanglement, nonclassical correlations, and nonlocal correlations are all forms of correlations between two or more subsystems of a composite quantum system that are different from strictly classical correlations and, in general, different from each other. One of the characteristic traits of classical correlations is that they can be freely shared. A party A can have maximal classical correlations with two parties B and C simultaneously. This is no longer the case if quantum entanglement or nonlocal correlations are concerned [1]. The limits on the shareability of those types of nonclassical correlations are known as monogamy constraints, see Fig. 1 for illustration. Strict monogamy inequalities have been proven that constrain the distribution of particular measures of entanglement and nonlocal correlations (the latter expressed in terms of violation of some Bell-type inequality [2]) among the subsystems of a multipartite system [3–11]. These relations can be seen as a particular case of trade-off relations that, in general, may relate and constrain different quantifiers of correlations [10,12]. Monogamy is the crucial property of correlations that makes quantum key distribution secure [1,13], even in no-signalling theories more general than quantum mechanics.

Nonclassical correlations that go beyond entanglement, often quantified, e.g., via the quantum discord [14,15], have recently attracted considerable attention [16,17]. While entanglement captures the nonseparability of two subsystems [18,19], quantum discord detects nonclassical properties even in separable states. Different attempts were presented to connect the new concept of quantum discord to quantum entanglement [20–26] and to broadcasting [27–29]. Several experimental results have been reported in [30–33]. Quantum discord, as well as related quantifiers of quantum correlations [17,22,23,34–44], have also been linked to better-than-classical performance in quantum

computation and communication tasks, even in the presence of limited or strictly vanishing entanglement [30,45–53]. An important question to understand the role of quantum correlations as signatures of genuine nonclassical behavior is whether they distribute in a monogamous way among multipartite systems.

A bipartite measure of correlations \mathcal{Q} satisfies monogamy if [3,19]

$$\mathcal{Q}^{A|BC}(\rho_{ABC}) \geq \mathcal{Q}^{A|B}(\rho_{AB}) + \mathcal{Q}^{A|C}(\rho_{AC}) \quad (1)$$

holds for all states ρ_{ABC} . Here, $\rho_{AB} = \text{Tr}_C(\rho_{ABC})$ denotes the reduced state of parties A and B , and analogously for ρ_{AC} . The vertical bar is the familiar notation for the bipartite split. The concept of monogamy is visualized in Fig. 1.




FIG. 1 (color online). Entanglement is monogamous: for a fixed amount of entanglement between A and BC , the more entanglement exists between A and B , the less can exist between A and C . Quantitatively, this is expressed using the monogamy relation, see Eq. (1) in the main text. In particular, the latter implies—for a monogamous measure of entanglement E —that $E^{A|C} = 0$ if $E^{A|BC} = E^{A|B}$. In this Letter we show that the monogamy relation does not hold, in general, for any quantum correlation measure beyond entanglement, i.e., for any measure that does not vanish on separable states.

If \mathcal{Q} denotes, in particular, an entanglement measure [18,19], then there are a number of choices that satisfy monogamy for pure states of qubits, including the squared concurrence [3] and the squared negativity [54], as well as their continuous variable counterparts for multimode Gaussian states [5,6]. The only known measure that is monogamous in all dimensions is the squashed entanglement [10,55]. Other entanglement measures, such as the entanglement of formation, do not satisfy the monogamy relation [3]. There is no known *a priori* rule about whether a given entanglement measure is monogamous or not. It is natural to ask whether a given measure for general quantum correlations is monogamous. Certain measures of general quantum correlations, such as quantum discord, were shown to violate monogamy by finding explicit examples of states for which the inequality (1) does not hold [56–61]. Those examples, however, do not exclude the possibility that other measures of quantum correlations, akin to the quantum discord, could exist that do satisfy a monogamy inequality.

In this Letter we address the issue of whether monogamy, in general, can extend to general quantum correlations beyond entanglement. Quantitatively, this question can be formulated as follows: Does there exist a measure of correlations \mathcal{Q} that obeys the monogamy relation (1) and is nonzero on a separable state? We will put this question to rest by proving that all measures for quantum correlations beyond entanglement (i.e., that are nonvanishing on at least some separable state) and that respect some basic properties are not monogamous in general. These basic properties of the correlation measure \mathcal{Q} are the following:

(1) positivity, i.e.,

$$\mathcal{Q}^{A|B}(\rho_{AB}) \geq 0; \quad (2)$$

(2) invariance under local unitaries $U_A \otimes V_B$, i.e.,

$$\mathcal{Q}^{A|B}(\rho_{AB}) = \mathcal{Q}^{A|B}(U_A \otimes V_B \rho_{AB} U_A^\dagger \otimes V_B^\dagger); \quad (3)$$

(3) no increase upon attaching a local pure ancilla, i.e.,

$$\mathcal{Q}^{A|B}(\rho_{AB}) \geq \mathcal{Q}^{A|BC}(\rho_{AB} \otimes |0\rangle\langle 0|_C). \quad (4)$$

These properties are valid for several measures of correlations known in the literature, including all entanglement measures [18,19]. In particular, positivity and invariance under local unitaries are standard requirements [62]. For the quantum discord defined in Refs. [14,15], which is an asymmetric quantity, Eq. (4) can be verified by inspection and is valid independently of whether the ancilla is attached on the side where the measurement entering the definition of discord is to be performed or on the unmeasured side. In a more general scenario, quantum correlations can be defined as the minimal distance to the set of classically correlated states [23,38,39,41]. In this case, Eq. (4) follows from the fact that any “reasonable” distance does not change upon attaching an ancilla: $D(\rho, \sigma) = D(\rho \otimes |0\rangle\langle 0|, \sigma \otimes |0\rangle\langle 0|)$. The same arguments can be

applied to measures that are defined via measurements on local subsystems [36]. Alternatively, quantum correlations may be investigated and quantified in terms of the minimal amount of entanglement necessarily created between the system and a measurement apparatus realizing a complete projective measurement [22,23,26,63]. Equation (4) also holds in this case, which can be seen solely using the properties of entanglement measures.

We are now in position to prove the following theorem.

Theorem 1. A measure of correlations \mathcal{Q} that respects Eqs. (2)–(4), and is also monogamous according to (1) must vanish for all separable states.

Proof.— Consider a measure \mathcal{Q} respecting the hypothesis and a generic separable state $\rho_{AC} = \sum_i p_i |\psi_i\rangle\langle\psi_i|_A \otimes |\phi_i\rangle\langle\phi_i|_C$. In the following, we will concentrate on a special extension of ρ_{AC} , defined as

$$\rho_{ABC} = \sum_i p_i |\psi_i\rangle\langle\psi_i|_A \otimes |i\rangle\langle i|_B \otimes |\phi_i\rangle\langle\phi_i|_C, \quad (5)$$

with orthogonal states $\{|i\rangle_B\}$. Observe that ρ_{ABC} has the same amount of correlations $\mathcal{Q}^{A|BC}$ as the state

$$\sigma_{ABC} = \sum_i p_i |\psi_i\rangle\langle\psi_i|_A \otimes |i\rangle\langle i|_B \otimes |0\rangle\langle 0|_C, \quad (6)$$

since both states are related by a local unitary on BC . On the other hand, Eq. (4) implies that σ_{ABC} does not have more correlations than the reduced state σ_{AB} . Taking these two observations together, we obtain $\mathcal{Q}^{A|B}(\sigma_{AB}) \geq \mathcal{Q}^{A|BC}(\rho_{ABC})$. Now, we invoke the monogamy relation for the state ρ_{ABC} , which leads us to the inequality

$$\mathcal{Q}^{A|B}(\sigma_{AB}) \geq \mathcal{Q}^{A|B}(\rho_{AB}) + \mathcal{Q}^{A|C}(\rho_{AC}). \quad (7)$$

The final ingredient in the proof is the fact that the two states ρ_{AB} and σ_{AB} are equal. From the positivity of the measure, it follows immediately that $\mathcal{Q}^{A|C}$ must vanish on the state ρ_{AC} . Since the latter is a generic separable state, \mathcal{Q} must vanish on all separable states. ■

The power of Theorem 1 lies in its generality. Under very weak assumptions, it rules out the existence of monogamous correlations beyond entanglement. Note that the arguments used in the proof of Theorem 1 are strong enough to show that the violation of monogamy appears even in three-qubit systems. This can be seen starting from Eq. (5), with each subsystem being a qubit. The measure \mathcal{Q} violates monogamy if it is nonzero on some separable two-qubit state of rank two. This is the case for quantum discord and any related measures of quantum correlations.

As we have argued below Eq. (4), the properties (2)–(4) are satisfied by all reasonable measures of quantum correlations known to the authors. However, in general, it cannot be excluded that the measure under study violates one of the properties given in Eqs. (2) and (3), or (4). Alternatively, we assume that some of these properties cannot be proven. In this situation, Theorem 1 does not tell us whether \mathcal{Q} is monogamous or not. Then, it is still possible to show that a monogamous measure \mathcal{Q} must be

zero on all separable states if it remains finite for a fixed dimension of one subsystem, i.e., if

$$\mathcal{Q}^{A|B} \leq f(d_A) < \infty \quad (8)$$

for fixed d_A , and some function f . To see this, we use the fact that any separable state ρ_{AB} has a symmetric extension $\rho_{AB_1 \dots B_n}$ such that $\rho_{AB} = \rho_{AB_i}$ holds for all $1 \leq i \leq n$, where n is an arbitrary positive integer [64–67]. Equation (8) implies that the measure $\mathcal{Q}^{A|B_1 \dots B_n}(\rho_{AB_1 \dots B_n})$ is finite for all n , including the limit $n \rightarrow \infty$. On the other hand, if \mathcal{Q} is monogamous, it has to fulfill the following inequality:

$$\mathcal{Q}^{A|B_1 \dots B_n}(\rho_{AB_1 \dots B_n}) \geq n \mathcal{Q}^{A|B}(\rho_{AB}). \quad (9)$$

However, if the measure \mathcal{Q} is nonzero on the separable state ρ_{AB} , one can always choose some n which is large enough such that Eq. (9) is violated, and thus \mathcal{Q} cannot be monogamous.

So far we have presented two different ways to show that a given measure of quantum correlations \mathcal{Q} violates monogamy, namely, Theorem 1 and Eq. (8). At this stage, it is natural to ask whether these two results have the same power, i.e., whether they allow us to draw the same conclusions about the structure of a given measure \mathcal{Q} . As already noted above, the proof of Theorem 1 allows us to rule out monogamy even for the simplest case of three qubits, as long as the measure \mathcal{Q} does not vanish on some separable state of two qubits having rank not larger than two. On the other hand, this argument does not apply to Eqs. (8) and (9). Indeed, if \mathcal{Q} is nonzero on some separable two-qubit state ρ_{AB} , Eqs. (8) and (9) only allow the statement that the measure \mathcal{Q} violates monogamy for some extension $\rho_{AB_1 \dots B_n}$. In particular, if $n > 2$, this result does not provide any insight about the monogamy of the measure for three-qubit states.

We move on to observe that monogamy [Eq. (1)], together with positivity [Eq. (2)], invariance under local unitary [Eq. (3)], and no increase under attaching a local ancilla [Eq. (4)] imply no increase under local operations. This is due to the fact that any quantum operation Λ admits a Stinespring dilation: $\Lambda[\rho_B] = \text{Tr}_C(U_{BC}\rho_B \otimes |0\rangle\langle 0|_C U_{BC}^\dagger)$; i.e., any quantum operation can be seen as resulting from a unitary operation on a larger-dimensional Hilbert space. Thus, for \mathcal{Q} respecting Eqs. (1)–(4), one finds

$$\begin{aligned} \mathcal{Q}^{A|B}(\rho_{AB}) &\geq \mathcal{Q}^{A|BC}(\rho_{AB} \otimes |0\rangle\langle 0|_C) \\ &= \mathcal{Q}^{A|BC}(U_{BC}\rho_{AB} \otimes |0\rangle\langle 0|_C U_{BC}^\dagger) \\ &\geq \mathcal{Q}^{A|B}(\text{Tr}_C(U_{BC}\rho_{AB} \otimes |0\rangle\langle 0|_C U_{BC}^\dagger)) \\ &\quad + \mathcal{Q}^{A|C}(\text{Tr}_B(U_{BC}\rho_{AB} \otimes |0\rangle\langle 0|_C U_{BC}^\dagger)) \\ &\geq \mathcal{Q}^{A|B}(\Lambda_B[\rho_{AB}]). \end{aligned} \quad (10)$$

No-increase under local operations [68], and thus, *a fortiori*, monogamy [the latter together with the almost trivial properties (2)–(4)] imply the following.

Theorem 2. A measure of correlations \mathcal{Q} that is non-increasing under operations on at least one side must be maximal on pure states; that is, for any ρ_{AB} on $\mathbb{C}^d \otimes \mathbb{C}^d$ there exists a pure state $|\psi\rangle\langle\psi|_{AB} \in \mathbb{C}^d \otimes \mathbb{C}^d$ such that $\mathcal{Q}^{A|B}(|\psi\rangle\langle\psi|_{AB}) \geq \mathcal{Q}^{A|B}(\rho_{AB})$.

Proof.—Immediate when one uses the fact that any state ρ_{AB} can be seen as the result of the application of a channel Λ_B (Λ_A) on any purification $|\psi\rangle_{AB}$ of ρ_A (ρ_B) (see, for example, [55]). Suppose that the measure \mathcal{Q} is nonincreasing under quantum operations on A . Then:

$$\mathcal{Q}^{A|B}(|\psi\rangle\langle\psi|_{AB}) \geq \mathcal{Q}^{A|B}(\Lambda_A[|\psi\rangle\langle\psi|_{AB}]) = \mathcal{Q}^{A|B}(\rho_{AB}). \quad (11)$$

■

This simple theorem is relevant, in particular, for the case of symmetric measures of quantum correlations. Several such measures were proposed in Refs. [23,38,41]. Some of these measures have counterintuitive properties. In particular, in [23] it was shown that for the relative entropy of quantumness, there exist mixed states ρ_{AB} that have more quantum correlations than any pure state $|\psi\rangle_{AB}$. The theorem just proven can be interpreted as a signature of the fact that general quantum correlations can increase under local operations (and *a fortiori* as a signature of the lack of monogamy) [41].

Theorem 1 and the reasoning in its proof amount essentially to the following insight about the violation of monogamy: if there is a separable state ρ_{AB} with nonzero correlations \mathcal{Q} , then there exists a mixed state ρ_{ABC} which proves that the measure under scrutiny is not monogamous: $\mathcal{Q}^{A|BC}(\rho_{ABC}) < \mathcal{Q}^{A|B}(\rho_{AB}) + \mathcal{Q}^{A|C}(\rho_{AC})$. On the other hand, crucially, a measure of correlations can still respect monogamy when evaluated on pure states $\rho_{ABC} = |\psi\rangle\langle\psi|_{ABC}$. As will be demonstrated in the following, the geometric measure of discord has exactly this property for three qubits. Before we present this result, we recall the definition of this measure.

The geometric measure of discord D_G was defined in Ref. [39] as the minimal square Hilbert-Schmidt distance to the set of classical-quantum states (CQ):

$$D_G^{A|B}(\rho_{AB}) = \min_{\sigma_{AB} \in \text{CQ}} \|\rho_{AB} - \sigma_{AB}\|_2^2. \quad (12)$$

Here, we used the 2-norm, also known as Hilbert-Schmidt norm, $\|\rho - \sigma\|_2 = \sqrt{\text{Tr}(\rho - \sigma)^2}$, and the minimum is taken over all classical-quantum states σ_{AB} . These are states which can be written as $\sigma_{AB} = \sum_i p_i |i\rangle\langle i|_A \otimes \sigma_B^i$ with some local orthogonal basis $\{|i\rangle_A\}$. The geometric discord has an operational interpretation in terms of the average fidelity of the remote state preparation protocol for two-qubit systems [69]. As noted above, the geometric measure of discord cannot be monogamous in general, since it is nonzero on some separable states. However, the following holds.

Theorem 3. The geometric measure of discord is monogamous for all pure states $|\psi\rangle_{ABC}$ of three qubits:

$$D_G^{ABC}(|\psi\rangle\langle\psi|_{ABC}) \geq D_G^{AB}(\rho_{AB}) + D_G^{AC}(\rho_{AC}), \quad (13)$$

where $\rho_{AB} = \text{Tr}_C(|\psi\rangle\langle\psi|_{ABC})$ and analogously for ρ_{AC} .

Proof.—We notice that for proving the inequality in Eq. (13), it is enough to show that for any pure state $|\psi\rangle_{ABC}$ there exists a classical-quantum state σ_{ABC} such that

$$D_G^{ABC}(|\psi\rangle\langle\psi|_{ABC}) \geq \|\rho_{AB} - \sigma_{AB}\|_2^2 + \|\rho_{AC} - \sigma_{AC}\|_2^2. \quad (14)$$

This inequality then automatically implies inequality (13), as, due to the minimization in the geometric measure of discord, the right-hand side of (13) can only be smaller than or equal to the right-hand side of (14). In order to show the existence of the mentioned classical-quantum state σ_{ABC} we choose a specific parametrization for a pure state of three qubits [70]:

$$\begin{aligned} |\psi_{ABC}\rangle = & \sqrt{p}|0\rangle_A(a|00\rangle_{BC} + \sqrt{1-a^2}|11\rangle_{BC}) \\ & + \sqrt{1-p}|1\rangle_A[\gamma(\sqrt{1-a^2}|00\rangle_{BC} - a|11\rangle_{BC}) \\ & + f|01\rangle_{BC} + g|10\rangle_{BC}]. \end{aligned} \quad (15)$$

The real numbers p , a , and f range between 0 and 1, g is complex with $0 \leq f^2 + |g|^2 \leq 1$, and $\gamma = \sqrt{1-f^2-|g|^2}$ is also real.

We proceed by evaluating the left-hand side of Eq. (14), using the explicit formula for pure states [71,72]:

$$D_G^{ABC}(|\psi\rangle\langle\psi|_{ABC}) = 2(1-p)p. \quad (16)$$

In the next step, we define the classical-quantum state $\sigma_{ABC} = \sum_{i=0}^1 \Pi_A^i \rho_{ABC} \Pi_A^i$ with local projectors in the computational basis: $\Pi_A^i = |i\rangle\langle i|_A$. The evaluation of the right-hand side of Eq. (14) is straightforward:

$$\|\rho_{AB} - \sigma_{AB}\|_2^2 + \|\rho_{AC} - \sigma_{AC}\|_2^2 = 2c(1-p)p \quad (17)$$

with $c = 1 + [4a^2(1-a^2) - 1]\gamma^2$. The proof is complete, if we can show that c cannot be larger than 1. This can be seen by noting that the term $4a^2(1-a^2)$ is maximal for $a^2 = \frac{1}{2}$, which leads to the maximal possible value $c = 1$. ■

Even though quantum correlations beyond entanglement cannot be monogamous in general, Theorem 3 demonstrates that for pure states of three qubits, monogamy of the geometric measure of discord is still preserved. To the best of our knowledge, this is the first instance of a measure of quantum correlations beyond entanglement that satisfies a restricted monogamy inequality. Certainly, this is not a property that all measures of quantum correlations have in common: As shown, e.g., in Ref. [56], the original quantum discord violates monogamy even on some pure states of three qubits.

In conclusion, we have addressed the question of monogamy for quantum correlations beyond entanglement. Using very general arguments, we have proven that any measure of correlations which is nonzero on some separable state unavoidably violates monogamy. Furthermore, we have shown that any monogamous measure of quantum correlations must be maximal on pure states. These results imply severe constraints on any monogamous measure of quantum correlations, and can also be used to witness the violation of monogamy. Finally, we have shown that even though all measures of nonclassical correlations akin to quantum discord cannot be monogamous for all states, they still may obey monogamy in certain restricted situations. In particular, we proved that the geometric measure of discord is monogamous for all pure states of three qubits. It is an open question whether there exists a measure of general quantum correlations which is monogamous for tripartite pure states of arbitrary dimensions. Another open question, which points to a possible future research direction, arises from the generalization of quantum discord to theories which are more general than quantum [73]. We hope that the results presented in this Letter are also useful for this more general scenario. Thus, the answer to the question posed in the title is: General quantum correlations are, in general, not monogamous.

We thank Davide Girolami and Hermann Kampermann for discussions. M. P. acknowledges support by NSERC, CIFAR, Ontario Centres of Excellence. G. A. is supported by a Nottingham Early Career Research and Knowledge Transfer Award. M. P. and G. A. acknowledge joint support by the EPSRC Research Development Fund (Pump Priming Grant No. 0312/09). G. A. acknowledges ESF for sponsoring the workshop during which this work was started. D. B. and A. S. acknowledge financial support by DFG; A. S. was supported by ELES.

-
- [1] B. M. Terhal, *IBM J. Res. Dev.* **48**, 71 (2004).
 - [2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
 - [3] V. Coffman, J. Kundu, and W. K. Wootters, *Phys. Rev. A* **61**, 052306 (2000).
 - [4] T. J. Osborne and F. Verstraete, *Phys. Rev. Lett.* **96**, 220503 (2006).
 - [5] G. Adesso and F. Illuminati, *New J. Phys.* **8**, 15 (2006).
 - [6] T. Hiroshima, G. Adesso, and F. Illuminati, *Phys. Rev. Lett.* **98**, 050503 (2007).
 - [7] B. Toner, *Proc. R. Soc. A* **465**, 59 (2009).
 - [8] J. Barrett, A. Kent, and S. Pironio, *Phys. Rev. Lett.* **97**, 170409 (2006).
 - [9] M. Pawłowski and Č. Brukner, *Phys. Rev. Lett.* **102**, 030403 (2009).
 - [10] M. Koashi and A. Winter, *Phys. Rev. A* **69**, 022309 (2004).
 - [11] M. P. Seevinck, *Quant. Info. Proc.* **9**, 273 (2010).
 - [12] M. Horodecki and M. Piani, *J. Phys. A* **45**, 105306 (2012).
 - [13] M. Pawłowski, *Phys. Rev. A* **82**, 032313 (2010).

- [14] H. Ollivier and W. H. Zurek, *Phys. Rev. Lett.* **88**, 017901 (2001).
- [15] L. Henderson and V. Vedral, *J. Phys. A* **34**, 6899 (2001).
- [16] Z. Merali, *Nature (London)* **474**, 24 (2011).
- [17] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, [arXiv:1112.6238](https://arxiv.org/abs/1112.6238).
- [18] M. B. Plenio and S. Virmani, *Quantum Inf. Comput.* **7**, 1 (2007).
- [19] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [20] V. Madhok and A. Datta, *Phys. Rev. A* **83**, 032323 (2011).
- [21] D. Cavalcanti, L. Aolita, S. Boixo, K. Modi, M. Piani, and A. Winter, *Phys. Rev. A* **83**, 032324 (2011).
- [22] A. Streltsov, H. Kampermann, and D. Bruß, *Phys. Rev. Lett.* **106**, 160401 (2011).
- [23] M. Piani, S. Gharibian, G. Adesso, J. Calsamiglia, P. Horodecki, and A. Winter, *Phys. Rev. Lett.* **106**, 220403 (2011).
- [24] M. F. Cornelio, M. C. de Oliveira, and F. F. Fanchini, *Phys. Rev. Lett.* **107**, 020502 (2011).
- [25] F. F. Fanchini, M. F. Cornelio, M. C. de Oliveira, and A. O. Caldeira, *Phys. Rev. A* **84**, 012313 (2011).
- [26] M. Piani and G. Adesso, *Phys. Rev. A* **85**, 040301(R) (2012).
- [27] M. Piani, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **100**, 090502 (2008).
- [28] M. Piani, M. Christandl, C. E. Mora, and P. Horodecki, *Phys. Rev. Lett.* **102**, 250503 (2009).
- [29] S. Luo and W. Sun, *Phys. Rev. A* **82**, 012338 (2010).
- [30] B. P. Lanyon, M. Barbieri, M. P. Almeida, and A. G. White, *Phys. Rev. Lett.* **101**, 200501 (2008).
- [31] J. Xu, X. Xu, C. Li, C. Zhang, X. Zou, and G. Guo, *Nature Commun.* **1**, 1 (2010).
- [32] R. Auccaise, J. Maziero, L. C. Céleri, D. O. Soares-Pinto, E. R. deAzevedo, T. J. Bonagamba, R. S. Sarthour, I. S. Oliveira, and R. M. Serra, *Phys. Rev. Lett.* **107**, 070501 (2011).
- [33] G. Passante, O. Moussa, D. A. Trottier, and R. Laflamme, *Phys. Rev. A* **84**, 044302 (2011).
- [34] J. Oppenheim, M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **89**, 180402 (2002).
- [35] A. K. Rajagopal and R. W. Rendell, *Phys. Rev. A* **66**, 022104 (2002).
- [36] S. Luo, *Phys. Rev. A* **77**, 022301 (2008).
- [37] S. Wu, U. V. Poulsen, and K. Mølmer, *Phys. Rev. A* **80**, 032319 (2009).
- [38] K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson, *Phys. Rev. Lett.* **104**, 080501 (2010).
- [39] B. Dakić, V. Vedral, and Č. Brukner, *Phys. Rev. Lett.* **105**, 190502 (2010).
- [40] R. Rossignoli, N. Canosa, and L. Ciliberti, *Phys. Rev. A* **82**, 052342 (2010).
- [41] A. Streltsov, H. Kampermann, and D. Bruß, *Phys. Rev. Lett.* **107**, 170502 (2011).
- [42] G. L. Giorgi, B. Bellomo, F. Galve, and R. Zambrini, *Phys. Rev. Lett.* **107**, 190501 (2011).
- [43] M. D. Lang, C. M. Caves, and A. Shaji, [arXiv:1105.4920](https://arxiv.org/abs/1105.4920).
- [44] J. Xu, *J. Phys. A* **44**, 445310 (2011).
- [45] E. Knill and R. Laflamme, *Phys. Rev. Lett.* **81**, 5672 (1998).
- [46] D. P. DiVincenzo, M. Horodecki, D. W. Leung, J. A. Smolin, and B. M. Terhal, *Phys. Rev. Lett.* **92**, 067902 (2004).
- [47] A. Datta, A. Shaji, and C. M. Caves, *Phys. Rev. Lett.* **100**, 050502 (2008).
- [48] A. Brodutch and D. R. Terno, *Phys. Rev. A* **83**, 010301 (2011).
- [49] R. Chaves and F. de Melo, *Phys. Rev. A* **84**, 022324 (2011).
- [50] B. Eastin, [arXiv:1006.4402](https://arxiv.org/abs/1006.4402).
- [51] K. Modi, H. Cable, M. Williamson, and V. Vedral, *Phys. Rev. X* **1**, 021022 (2011).
- [52] V. Madhok and A. Datta, [arXiv:1107.0994](https://arxiv.org/abs/1107.0994).
- [53] S. Boixo, L. Aolita, D. Cavalcanti, K. Modi, and A. Winter, *Int. J. Quantum. Inform.* **09**, 1643 (2011).
- [54] Y.-C. Ou and H. Fan, *Phys. Rev. A* **75**, 062308 (2007).
- [55] M. Christandl and A. Winter, *J. Math. Phys. (N.Y.)* **45**, 829 (2004).
- [56] G. L. Giorgi, *Phys. Rev. A* **84**, 054301 (2011).
- [57] R. Prabhu, A. K. Pati, A. S. De, and U. Sen, [arXiv:1109.4318](https://arxiv.org/abs/1109.4318).
- [58] R. Prabhu, A. K. Pati, A. S. De, and U. Sen, *Phys. Rev. A* **85**, 040102(R) (2012).
- [59] Sudha, A. R. Usha Devi, and A. K. Rajagopal, [arXiv:1110.3026](https://arxiv.org/abs/1110.3026).
- [60] M. Allegra, P. Giorda, and A. Montorsi, *Phys. Rev. B* **84**, 245133 (2011).
- [61] X.-J. Ren and H. Fan, [arXiv:1111.5163](https://arxiv.org/abs/1111.5163).
- [62] A. Brodutch and K. Modi, [arXiv:1108.3649](https://arxiv.org/abs/1108.3649).
- [63] S. Gharibian, M. Piani, G. Adesso, J. Calsamiglia, and P. Horodecki, *Int. J. Quantum. Inform.* **09**, 1701 (2011).
- [64] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).
- [65] R. F. Werner, *Lett. Math. Phys.* **17**, 359 (1989).
- [66] A. C. Doherty, P. A. Parrilo, and F. M. Spedalieri, *Phys. Rev. A* **69**, 022308 (2004).
- [67] D. Yang, *Phys. Lett. A* **360**, 249 (2006).
- [68] To be precise, no-increase just under operations on one side.
- [69] B. Dakić, Y. O. Lipp, X. Ma, M. Ringbauer, S. Kropatschek, S. Barz, T. Paterek, V. Vedral, A. Zeilinger, Č. Brukner, and P. Walther, [arXiv:1203.1629](https://arxiv.org/abs/1203.1629).
- [70] T. A. Brun and O. Cohen, *Phys. Lett. A* **281**, 88 (2001).
- [71] S. Luo and S. Fu, *Phys. Rev. Lett.* **106**, 120401 (2011).
- [72] D. Girolami and G. Adesso, *Phys. Rev. A* **84**, 052110 (2011).
- [73] P. Perinotti, *Phys. Rev. Lett.* **108**, 120502 (2012).